



MaxGrow®

0-0-0 + 1% (Zn)

GUARANTEED ANALYSIS: ZINC (Zn).....1.0%

THIS PRODUCT IS DERIVED FROM ZINC SULFATE

CAUTION: KEEP OUT OF REACH OF CHILDREN



WARNING

Harmful if swallowed, Causes skin irritation, Causes eye irritation

PRECAUTIONARY STATEMENTS:

Avoid breathing fume/mist/vapors/spray. Wash skin thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well ventilated area. Wear protective gloves/protective clothing/eye and face protection.

IF SWALLOWED: Call a poison control or doctor/physician if you feel unwell. Rinse mouth.

IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Call a poison control center or doctor/physician if you feel unwell.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

Take off contaminated clothing and wash it before reuse. Store in a well ventilated place. Keep container tightly closed. Store locked up. Dispose of contents/container to an approved waste disposal plant in accordance with local/regional/national regulations.

Purpose: MaxGrow is a liquid nutrient product that is designed to provide Zinc (Zn) that may be deficient in the soil or plant.

GENERAL DIRECTIONS FOR USE:

Use 4 to 25 oz. / acre. Apply by water run (flood or sprinkler), spray (air or ground), injected into fertigation system, or shank in with or without nutrient. Although compatible with most pesticides and nutrients, a jar test is recommended. Otherwise, consult your technical representative

Shake well before using

STORAGE AND HANDLING:

Store in a sealed container in a cool dry place, out of direct sunlight. Store at temperatures above 40 degrees F and below 95 degrees F. Product must be used within 15 days after opening for maximum efficacy. Use within 36 hours of mixing.

Conditions of Sale and Warranty

1. Seller warrants that this material conforms to the description on the label and is reasonably fit for use as directed hereon. Seller neither makes nor authorizes any agent or representative to make any other warranty of fitness or of merchantability, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond seller's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon, and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by seller under 1. above) resulting from handling, storage and use of material

Information regarding the contents and levels of metal in this product is available on the internet at <http://www.aapfco.org/metals.htm>

SUGGESTED APPLICATION

Small Grains: 12.8 fl. oz./acre spray to soil at planting or near dormancy break and up to 25 fl. oz./acre foliar applied assists with plant vigor.

Row Crops: Corn, Soybeans and Sunflowers etc.: Spray to soil at or near planting on surface, in furrow or side banded. May also be used with side dressed plant food. Crop may benefit from 6-8 fl. oz./acre foliar spray to plant every 30 days.

Alfalfa: 12.8 fl. oz./acre spray to soil at or near dormancy break. May also apply 8 fl.oz./acre 7-10 days after each cutting throughout the season.

Vegetable Crops: 12.8 fl. oz./acre spray to soil at or near planting. Follow with 4-6 fl. oz./acre spray on plant every 30 days.

Nut Tree Crops: 25 fl. oz./acre spray to soil post harvest or early spring.

Vines and Canes: 25 fl. oz./acre spray to soil in fall or early spring.

Citrus: 25 fl. oz./acre twice per year spray to soil or through drip irrigation.

Turf: 12.8 fl. oz./acre to soil at planting followed by 12.8 fl. oz./acre every 34-40 days.

Nurseries: Apply a 1-3% solution of MaxGrow (1-3 gallons/100 gallons of water) through overhead sprinklers or drip system at each irrigation.



Weight per gallon: 8.56 (lb.) 3.88 (kg) at 68 degrees F.

Net Contents: 2.5 gallons (9.463 liter) Net Weight: 21.4 lbs. (9.70 kg)
Net Contents: 5 gallons (18.927 liter) Net Weight: 42.8 lbs. (19.41 kg)
Net Contents: 270 gallons (1022.06 liter) Net Weight: 2311 lbs. (1048 kg)